


SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)
Subject with Code: NUMERICAL METHODS AND TRANSFORMS (19HS0834)

Branch: B.Tech. (ECE)

Year & Sem: II-B.Tech. & I-Sem. **Regulation:** R19

UNIT –I

1. Find out the square root of 25 given $x_0 = 2.0$, $x_1 = 7.0$ using Bisection method. [12M]
2. Find a positive root of $x^3 - x - 1 = 0$ correct to two decimal places by Bisection method. [12M]
3. Find a positive root of $f(x) = e^x - 3$ correct to two decimal places by Bisection method. [12M]
4. Find a real root of the equation $xe^x - \cos x = 0$ using Newton – Raphson method. [12M]
5. Using Newton-Raphson method (i) Find square root of 28 (ii) Find cube root of 15. [12M]
6. a) Using Newton-Raphson method Find reciprocal of 12. [6M]
 b) Find a real root of the equation $x \tan x + 1 = 0$ using Newton – Raphson method. [6M]
7. Find out the root of the equation $x \log_{10}(x) = 1.2$ using False position method. [12M]
8. Find the root of the equation $xe^x = 2$ using Regula-falsi method. [12M]
9. From the following table values of x and $y = \tan x$. Interpolate values of y when $x = 0.12$ and $x = 0.28$.

x	0.10	0.15	0.20	0.25	0.30
y	0.1003	0.1511	0.2027	0.2553	0.3093
10. a) Using Newton's forward interpolation formula and the given table of values

x	1.1	1.3	1.5	1.7	1.9
$f(x)$	0.21	0.69	1.25	1.89	2.61

 Obtain the value of $f(x)$ when $x = 1.4$. [6M]
 b) Use Newton's backward interpolation formula to find $f(32)$ given $f(25) = 0.2707$, $f(30) = 0.3027$, $f(35) = 0.3386$, $f(40) = 0.3794$. [6M]



SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR
Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)

Subject with Code: NUMERICAL METHODS AND TRANSFORMS (19HS0834)

Branch: B.Tech. (ECE)

Year & Sem: II-B.Tech. & I-Sem. **Regulation:** R19

UNIT –II

1. Tabulate $y(0.1)$, $y(0.2)$ and $y(0.3)$ using Taylor's series method
given that $y' = y^2 + x$ and $y(0) = 1$ [12M]
2. Using Taylor's series method find an approximate value of y at $x = 0.2$ for the
D.E $y' - 2y = 3e^x$, $y(0) = 0$. Compare the numerical solution obtained with exact solution. [12M]
3. a) Solve $y' = x + y$, given $y(1) = 0$ find $y(1.1)$ and $y(1.2)$ by Taylor's series method. [6M]
b) Solve by Euler's method $\frac{dy}{dx} = \frac{2y}{x}$ given $y(1) = 2$ and find $y(2)$ [6M]
4. Using Euler's method, find an approximate value of y corresponding to $x = 1$ given that
 $\frac{dy}{dx} = x + y$ and $y = 1$ when $x = 0$ taking step size $h = 0.1$ [12M]
5. a) Using Euler's method $y' = y^2 + x$, $y(0) = 1$. Find $y(0.1)$ and $y(0.2)$ [6M]
b) Using Runge – Kutta method of fourth order, compute $y(0.2)$ from $y' = xy$, $y(0) = 1$,
taking $h = 0.2$ [6M]
6. Using R-K method of 4th order, solve $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}$, $y(0) = 1$. Find $y(0.2)$ and $y(0.4)$. [12M]
7. Using R-K method of 4th order find $y(0.1)$, $y(0.2)$ and $y(0.3)$ given that $\frac{dy}{dx} = 1 + xy$, $y(0) = 2$. [12M]
8. Solve $y'' - x(y')^2 + y^2 = 0$ using R-K method of 4th order for $x = 0.2$ given $y(0) = 1$,
and $y'(0) = 0$ taking $h = 0.2$ [12M]
9. Evaluate $\int_0^1 \frac{1}{1+x} dx$ (i) by Trapezoidal rule and Simpson's $\frac{1}{3}$ rule.
(ii) using Simpson's $\frac{3}{8}$ rule and compare the result with actual value. [12M]
10. a) Compute $\int_0^4 e^x dx$ by Simpson's $\frac{3}{8}$ rule with 12 sub divisions. [6M]
b) Compute $\int_3^7 x^2 \log x dx$ using Trapezoidal rule and Simpson's $\frac{1}{3}$ rule by taking 10 sub divisions. [6M]


SIDDARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)
Subject with Code: NUMERICAL METHODS AND TRANSFORMS (19HS0834)

Branch: B.Tech. (ECE)

Year & Sem: II-B.Tech. & I-Sem. **Regulation:** R19

UNIT-III

1. a) Find the Laplace transform of $f(t) = e^{3t} - 2e^{-2t} + \sin 2t + \cos 3t + \sinh 3t - 2\cosh 4t + 9$. [6M]
 b) Find the Laplace transform of $f(t) = \cosh at \sin bt$. [6M]
2. a) Find the Laplace transform of $f(t) = \left(\sqrt{t} + \frac{1}{\sqrt{t}}\right)^3$. [6M]
 b) Find the Laplace transform of $f(t) = e^{4t} \sin 2t \cos t$. [6M]
3. a) Find the Laplace transform of $f(t) = t^2 e^{2t} \sin 3t$. [6M]
 b) Find the Laplace transform of $f(t) = \frac{1 - \cos at}{t}$. [6M]
4. a) Find the Laplace transform of $f(t) = \int_0^t e^{-t} \cos t \, dt$. [6M]
 b) Find the Laplace transform of $f(t) = e^{-4t} \int_0^t \frac{\sin 3t}{t} \, dt$. [6M]
5. a) Show that $\int_0^\infty t^2 e^{-4t} \cdot \sin 2t \, dt = \frac{11}{500}$, Using Laplace transform. [6M]
 b) Using Laplace transform, evaluate $\int_0^\infty \frac{\cos at - \cos bt}{t} \, dt$. [6M]
6. a) Find $L^{-1}\left\{\frac{3s-2}{s^2-4s+20}\right\}$ by using first shifting theorem. [6M]
 b) Find $L^{-1}\left\{\log\left(\frac{s-a}{s-b}\right)\right\}$ [6M]
7. a) Find $L^{-1}\left\{\frac{1}{(s^2+5)^2}\right\}$, using Convolution theorem. [6M]
 b) Find $L^{-1}\left\{\frac{s^2}{(s^2+4)(s^2+25)}\right\}$, using Convolution theorem. [6M]
8. a) Find the Inverse Laplace transform of $\frac{1}{s(s^2+a^2)}$ [6M]
 b) Find $L^{-1}\left\{s \log\left(\frac{s-1}{s+1}\right)\right\}$ [6M]
9. Using Laplace transform method to solve $y^{11} - 3y^1 + 2y = 4t + e^{3t}$ where $y(0) = 1, y^1(0) = 1$ [12M]
10. Solve the D.E. $\frac{d^2x}{dt^2} + 2\frac{dx}{dt} + x = 3te^{-t}$ using Laplace Transform given that
 $x(0) = 4; \frac{dx}{dt} = 0 \text{ at } t = 0$ [12M]


SIDDARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)
Subject with Code: NUMERICAL METHODS AND TRANSFORMS (19HS0834)

Branch: B.Tech. (ECE)

Year & Sem: II-B.Tech. & I-Sem. **Regulation:** R19

UNIT – IV

 1. a) Obtain the Fourier series expansion of $f(x) = x^2$ in the interval $(0, 2\pi)$. [6M]

 b) Obtain the Fourier series expansion of $f(x) = (x - x^2)$ in the interval $[-\pi, \pi]$. Hence show that

$$\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots = \frac{\pi^2}{12}. \quad [6M]$$

 2. a) Obtain the Fourier series expansion of $f(x) = (\pi - x)^2$ in $0 < x < 2\pi$ and deduce that

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots = \frac{\pi^2}{6}. \quad [6M]$$

 b) Find the Fourier series for the function $f(x) = x$ in $-\pi < x < \pi$. [6M]

 3. Find a Fourier series to represent the function $f(x) = e^x$ for $-\pi < x < \pi$. And hence derive a series for $\frac{\pi}{\sinh \pi}$. [12M]

 4. Find the Fourier series to represent the function $f(x) = x^2$ for $-\pi < x < \pi$ and hence show that

$$(i) \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots = \frac{\pi^2}{12}. \quad (ii) \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots = \frac{\pi^2}{6}.$$

$$(iii) \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8} \quad [12M]$$

 5. a) If $f(x) = |\sin x|$, expand $f(x)$ as a Fourier series in the interval $(-\pi, \pi)$ [6M]

 b) Find the half range cosine series for $f(x) = x$ in the interval $0 \leq x \leq \pi$. [6M]

 6. Expand the function $f(x) = |x|$ in $-\pi < x < \pi$ as a Fourier series and deduce that

$$\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8} \quad [12M]$$

 7. Find the half range sine series for $f(x) = x(\pi - x)$ in the interval $0 \leq x \leq \pi$ and deduce that

$$\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \dots = \frac{\pi^3}{32}. \quad [12M]$$

 8. a) Expand $f(x) = e^{-x}$ as a Fourier series in the interval $(-1, 1)$. [6M]

 b) Expand $f(x) = |x|$ as a Fourier series in the interval $(-2, 2)$. [6M]

 9. a) Find the half range sine series expansion of $f(x) = x^2$ when $0 < x < 4$. [6M]

 b) Find the half range cosine series expansion of $f(x) = x(2 - x)$ in $0 \leq x \leq 2$. [6M]

 10. Find half range Fourier cosine series of $f(x) = (x - 1)^2$ in $0 < x < 1$.

$$\text{Hence show that} \quad i) \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots = \frac{\pi^2}{6} \quad ii) \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12}. \quad [12M]$$


SIDDARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)
Subject with Code: NUMERICAL METHODS AND TRANSFORMS (19HS0834)

Branch: B.Tech. (ECE)

Year & Sem: II-B.Tech. & I-Sem. **Regulation:** R19

UNIT – V

1. Find the Fourier transform of $f(x) = \begin{cases} 1; |x| < a \\ 0; |x| > a \end{cases}$ and hence evaluate

$$i) \int_{-\infty}^{\infty} \frac{\sin p \cos p x}{p} dp \quad ii) \int_{-\infty}^{\infty} \frac{\sin p}{p} dp \quad iii) \int_0^{\infty} \frac{\sin p}{p} dp. \quad [12M]$$

2. Find the Fourier transform of $f(x) = \begin{cases} a^2 - x^2, |x| < a \\ 0, |x| > a > 0 \end{cases}$ Hence show that $\int_0^{\infty} \frac{\sin x - x \cos x}{x^3} dx = \frac{\pi}{4}$. [12M]

3. a) Find the Fourier transform of $f(x) = e^{-\frac{x^2}{2}}, -\infty < x < \infty$ [6M]

- b) If $F(p)$ is the complex Fourier transform of $f(x)$, then prove that the complex Fourier transform of $f(x) = \cos ax$ is $\frac{1}{2}[F(p+a) + F(p-a)]$ [6M]

4. a) Find the Fourier cosine transform of $f(x)$ defined by $f(x) = \begin{cases} \cos x & ; 0 < x < a \\ 0 & ; x \geq a \end{cases}$ [6M]

- b) If $F(P)$ is the complex Fourier transform of $f(x)$, then prove that the complex Fourier transform of $F\{f(x-a)\} = e^{ipa} \cdot F(P)$ [6M]

5. Find the Fourier sine and cosine transforms of $f(x) = \frac{e^{-ax}}{x}$ and deduce that

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} \sin sx \, dx = \tan^{-1}\left(\frac{s}{a}\right) - \tan^{-1}\left(\frac{s}{b}\right). \quad [12M]$$

6. Find the Fourier sine and cosine transforms of $f(x) = e^{-ax}, a > 0$ and hence deduce the integrals [12M]

$$(i) \int_0^{\infty} \frac{p \sin px}{a^2 + p^2} dp \quad (ii) \int_0^{\infty} \frac{\cos px}{a^2 + p^2} dp$$

7. a) Prove that $F[x^n f(x)] = (-i)^n \frac{d^n}{dp^n} [F(p)]$ [6M]

- b) Prove that $F_s\{x f(x)\} = -\frac{d}{dp} [F_c(p)]$ [6M]

8. a) Find the Fourier cosine transform of $e^{-ax} \cos ax, a > 0$ [6M]

- b) Find the Fourier cosine transform of $f(x) = \begin{cases} x, & \text{for } 0 < x < 1 \\ 2-x, & \text{for } 1 < x < 2 \\ 0, & \text{for } x > 2 \end{cases}$ [6M]

9. Find the finite Fourier sine and cosine transform of $f(x)$ defined by $f(x) = 2x$ where $0 < x < 2\pi$. [12M]

10. a) Find the finite Fourier sine transform of $f(x)$, defined by $f(x) = \begin{cases} x, & 0 \leq x \leq \frac{\pi}{2} \\ \pi - x, & \frac{\pi}{2} \leq x \leq \pi \end{cases}$ [6M]

- b) Find the inverse finite Fourier sine transform of $f(x)$, If $F_s(n) = \frac{16(-1)^{n-1}}{n^3}$, where n is a positive integer and $0 < x < 8$. [6M]